

Inpatient Antepartum, Labor and Delivery Guidelines
COVID 19 (PUI or CONFIRMED)
Labor management

- **Patient attire/location**
 - Patient to be met by RNs in PPE in Adult Lobby and pt & support will be masked
 - Patient to wear clean patient gown and surgical mask (during entire labor)
 - Restricted to the room at all times
 - Can use telemonitors and tub as needed
 - Isolated room (not necessarily neg pressure room) with appropriate Respiratory Illness Precautions sign
- **Support/caregiver**
 - Encouraged to wear protective gear (CONTACT & DROPLET) at all times. Required to at least wear surgical face mask at all times
 - Restricted to patient's room, not to enter common spaces including cafeteria
 - Uses patient bathroom
 - Must be asymptomatic and cannot have been co-habiting with the patient prior to admission
 - Handwashing often
 - If develops respiratory symptoms (cough, fevers, etc) must leave immediately
- **Anesthesia Considerations and Oxygen use**
 - **Strongly** encourage epidural use to avoid intubation in case emergency CS required
 - No nitrous oxide or remifentanyl PCA
 - Only nasal cannula use for maternal hypoxia (Oxygen is **NO LONGER** used for intrauterine resuscitation)
- **Labor Room Materials**
 - Ensure labor management supply box is in the room with limited materials (Labeled **Covid+/PUI Supply Box**)
 - Box includes everything from usual supply cart in labor room including vacuums
 - Ensure FSE and IUPC cords are in the room
 - If ultrasound used, needs to be sterilized per protocol
- **Monitoring**
 - VS and continuous pulse oximetry, individualized per orders
 - Strict I/Os
- **Safety huddles with team**
 - Team check in q2hrs: (may not require in-person physical exam each time)
 - Team to include: Bedside nurse, MFM attg, MD attg, Charge nurse, OB anesthesia, ICU PRN
 - Attention to Maternal Early Warning Criteria for sepsis
 - Review patient's clinical status, early intervention, low threshold for cesarean
- **Delivery Personnel**
 - Recommend MD attending only with Chief resident PRN
 - Bedside nurse & LDR Resource nurse
 - Pediatrics team (essential members only)
- **Delivery Team Donning and Doffing**
 - Delivery team (MDs & nurses) to wear NOVEL RESPIRATORY PPE (including N-95 & face shield) during 2nd stage. Don sterile delivery gloves OVER non-sterile blue gloves
 - Hand hygiene, doffing of gown/gloves in pt rm (below the neck) and the doffing of masks and eyewear (above the neck) outside pt room, hand hygiene again
- **Delivery Positioning and Specifics**
 - Recommend pt in stirrups to have control of delivery and avoid sitting on bed to avoid/reduce risk of fluid exposure.
 - Skin-to-skin and delayed cord clamping to be discussed on case-to-case basis with ICN depending on plans for co-localization vs 14-day separation
- **Pediatrics Team**
 - Limited personnel/essential members only
 - All must be in PPE including provider to receive baby (will wear N-95 and face shields due to potential need for neonatal intubation)
 - If resuscitation is required, stays in LDR, then transfers baby in isolette to ICN
- **Post-delivery Management**
 - Debrief to be completed by MD attg with bedside RN – dispo to stay in LDR vs ICU
 - LDR to be CLOSED until appropriate high cleaning completed (up to 2 hours)
 - Prior to transfer: aim to clear hallways, minimize contact between patient and other staff/ surfaces.
 - Pt to wear surgical mask and covered from neck down in clean sheet for transport to PP; return wheelchair/gurney to dirty LDR

Inpatient Antepartum, Labor and Delivery Guidelines
COVID 19 (PUI or CONFIRMED)
Antepartum

- **Notify OB Anesthesia & ICN if patient develops respiratory symptoms or is admitted as PUI/COVID+**
 - MFM attg and ICN attending to discuss where delivery to take place if needed (minimize contamination of multiple spaces and PPE used)
- **Patient attire/location**
 - Patient to wear clean gown and surgical mask
 - Restricted to room at all times (not to use shared spaces)
 - Isolated room
- **Support/caregiver**
 - Recommend protective gear (CONTACT & DROPLET) at all times. Required to at least wear surgical face mask at all times
 - Restricted to patient's room, not to enter common spaces including cafeteria
 - Uses patient bathroom
 - Must be asymptomatic and cannot have been co-habiting with the patient prior to admission
 - Handwashing often
 - If develops respiratory symptoms (cough, fevers, etc) must leave immediately
- **Additional materials**
 - If ultrasound used, needs to be sterilized, so limit to only necessary use that will directly change patient management
 - Use PDC (see algorithm) for US cleaning
 - Use only disposable stethoscopes
 - Discuss opening OR3 surgical set on case-by-case basis
- **Rounding/Direct Patient Care**
 - MFM Attending only. Other team members can use tele-health or phone
 - Limit additional physical exams to one provider if possible when indicated
- **Safety huddles with team**
 - Team check in q2hrs, continuous pulse ox, strict I/Os (may not require in-person physical exam each time)
 - Attention to Maternal Early Warning Criteria for sepsis
 - Includes bedside nurse, MFM attg, MD attg, Charge nurse, OB anesthesia, ICU PRN
 - Review patient's clinical status, early intervention, low threshold for cesarean. Consider sending D-dimer and other labs to better understand pt's individual clinical risk factors for poor COVID outcomes
- **Equipment**
 - Small amount of gloves, gel, FSE, amniohook in room
 - Collected in clear plastic box (Antepartum Med room)
- **Transport of patient**
 - Complete imaging studies in pt room as much as possible to avoid transport
 - If pt needs to be transported, must wear surgical mask and new gown. Drape clean sheet over patient
 - Minimize transport team. Transporter must wear CONTACT & DROPLET PPE during transport.
 - Don and doff per transport protocols (whenever going into and out of rooms/clinical spaces)
- **If antepartum patient goes into labor or is induced, will stay in room, will NOT move to labor room**
- **Medication considerations:**
 - **Betamethasone** per protocol in pt <34 weeks if respiratory symptoms are mild/moderate. For pts with severe symptoms, pts who have already had a course and pts >34 weeks, discuss with MFM
 - **Magnesium sulfate** use should be discussed with MFM depending on the clinical situation and risk of further respiratory compromise

Inpatient Antepartum, Labor and Delivery Guidelines
COVID 19 (PUI or CONFIRMED)
PLANNED/URGENT/EMERGENCY Cesarean Section Overview Guidelines

- **Decision making**
 - Make early decisions in controlled environment as much as possible (ie. Early diagnosis of arrest of descent, fetal intolerance)
 - Minimize need for emergency delivery
 - Maximize time we have to use approved PRECAUTIONS to minimize exposure to personnel
- **Pre-operative anesthesia**
 - Patient should be encouraged to have an early epidural as to decrease risk of needing emergency intubation
 - Plan CSE for non-urgent cesarean
- **Caregiver**
 - Only present in OR if urgent c-section (NOT if emergency c-section)
 - Wears PPE the entire time (CONTACT & DROPLET)
- **Communication with unit**
 - HUSC will activate **COVID batch page "91119"**
 - Charge nurse will utilize HUSC to advise people to CLEAR THE HALLWAY
 - Only essential team member involvement
 - Observer/Recorder to monitor personnel (donning and doffing of PPE)
 - OR RN Resource communicates when additional personnel needed in OR via PHONE to designated OUT OF OR person (ie if blood is needed, additional supplies)
 - Clear communication with PCA and Anesthesia tech (outside of OR)
- **Provider PPE use**
 - IF in OR during surgery, needs to be Novel Respiratory (N-95) for COVID +/-PUI/unknown or pending
 - Outside of patient room (droplet/contact)
 - Outside of ORs in sterile core (droplet/contact)
- **Surgical personnel**
 - See separate personnel log and document
- **Newborn management**
 - OB team to hand off baby to Peds receiving person with appropriate delayed cord clamping
 - Baby will be resuscitated in the room per usual workflow
 - No skin-to-skin
- **Donning and Doffing of PPE by surgical personnel**
 - Occurs OUTSIDE of OR with separate trash bin (sterile core)
 - Should don NEW PPE prior to transporting patient
- **Surgical Timeouts**
 - Should be performed by MD OB attg to determine urgency of case
 - Post-op Debrief should include ICU attending, MFM attending (phone)
 - All non-essential providers should leave the room for extubation. Return to the room after 15 mins have passed
- **Patient recovery**
 - Recovery in her intrapartum LDR isolation room vs ICU per anesthesia and OB team determined after surgery
 - Portable monitor set up in isolation LDR
- **OR clean up**
 - HUSC/Charge RN communicates with housekeeping staff of COVID positive OR status
 - High risk clean up (up to two hours)

Inpatient Antepartum, Labor and Delivery Guidelines
COVID 19 (PUI or CONFIRMED)
Postpartum/Discharge

- **Notify OB Anesthesia & ICN if patient develops respiratory symptoms or is admitted as PUI/COVID+**
 - Add patient to the Complicated postpartum list
- **Post-delivery Management**
 - Debrief to be completed by MD attg with bedside RN – dispo to stay in LDR vs ICU
 - LDR to be CLOSED until appropriate high cleaning completed (up to 2 hours)
 - Prior to transfer: aim to clear hallways, minimize contact between patient and other staff/ surfaces
 - Pt to wear surgical mask and covered from neck down in clean sheet for transport to PP; return wheelchair/gurney to dirty LDR
- **Try to discharge stable PUI patients home with neonate ASAP**
 - Add pt's MRN to COVID follow up list
- **Patient attire/location**
 - Patient to wear clean gown and surgical mask
 - Restricted to patient room (not to enter shared common spaces)
 - Isolated room
- **Support/caregiver**
 - Cannot have had prolonged unprotected contact with the patient prior to hospitalization (ie. has not been co-habiting with the patient)
 - Recommend protective gear (CONTACT & DROPLET) at all times
 - Required to at least wear surgical face mask at all times
 - Restricted to patient's room, not to enter common spaces including cafeteria
 - Uses patient bathroom
 - Must be asymptomatic and cannot have been co-habiting with the patient prior to admission
 - Handwashing often
 - If develops infectious symptoms (cough, fevers, etc) must leave immediately
- **Additional materials**
 - Use only disposable stethoscopes
- **Safety huddles with team**
 - Team check in q2hrs, continuous pulse ox, strict I/Os (may not require in-person physical exam each time)
 - Attention to Maternal Early Warning Criteria for sepsis
 - Includes bedside nurse, MFM attg, MD attg, Charge nurse, OB anesthesia, ICU PRN
- **Rounding/Direct Patient Care**
 - OB attending only
 - Other team members can use tele-health or phone
- **Equipment**
 - Small amount of materials in the room
- **Transport of patient**
 - Complete imaging studies in pt room as much as possible to avoid transport
 - If pt needs to be transported, must wear surgical mask and new gown. Drape new sheet over patient
 - Minimize personnel in transport team. Transporter must wear contact & droplet PPE during transport
 - Don and doff per transport protocols (whenever going into and out of rooms/clinical spaces)
- **Breastfeeding considerations**
 - Good hand hygiene and wear mask while feeding baby
 - Recommend to express and pump breast milk for infant if separated
 - Follow CDC guidelines for proper breast pumping hygiene
- **If patient requires prolonged hospital stay, consider transfer to Parnassus/Mt Zion to be managed by COVID medicine services**

Inpatient Antepartum, Labor and Delivery Guidelines
COVID 19 (PUI or CONFIRMED)
Safety huddle protocols and guidance

- **Complete safety huddles every 2 hours for ALL PUI/COVID + patients on the unit**
 - Assess labor process, fetal status, maternal status
 - Triage, intrapartum, antepartum, postpartum
 - Direct physical exam MAY NOT be indicated each huddle
- **Safety huddle personnel (outside patient room)**
 - Charge Nurse
 - Bedside nurse (1:1)
 - OB/MFM attending
 - OB anesthesia
 - Chief resident
 - ICU attending PRN
- **Monitoring for patients**
 - Continuous pulse oximetry
 - Strict I/Os
 - Attention to respiratory rate
 - LIMIT Physical exams to ONLY if there are vital sign abnormalities or change in clinical status

Questions to consider for Huddle:

- Maternal hemodynamic status (see below from the National Partnership for Maternal Safety)
 - Is the patient on oxygen support/tachypnea/decreased urine output/hypotension? → if YES, prompt evaluation of maternal status and reconsider proceeding with labor if remote from delivery
 - Discussion about indications for further testing → repeat CXR, CT, Labs
- Fetal Monitoring Category I? → YES/NO → if NO, low threshold to assess status and consider a controlled Cesarean section
- Assessment of labor progress → if in the active phase (greater than 6 cm) with no cervical change over 2 hours → consider interventions versus Cesarean delivery

Maternal Early Warning Criteria From the National Partnership for Maternal Safety

- Systolic BP (mm Hg) <90 or >160
- Diastolic BP (mm Hg) >100
- Heart rate (beats per min) <50 or >120
- Respiratory rate (breaths per min) <10 or >30
- Oxygen saturation on room air, at sea level, % <95
- Oliguria, mL/hr for ≥ 2 hrs <35
- Maternal agitation, confusion, or unresponsiveness; Patient with preeclampsia reporting a non-remitting headache or shortness of breath

Inpatient Antepartum, Labor and Delivery Guidelines

Changes in L&D management of the Asymptomatic patient

- **Testing guidelines**
 - All patients will be tested for COVID prior to admission for any planned procedure or admission (within 4 days of admission)
 - All patients without a COVID result in the past 4 days will be tested for COVID testing once admitted
 - If patient has been admitted for >7 days and has a high risk of requiring an aerosol generating procedure (AGP), she will be re-tested for COVID (ie. viable pregnancies)
- **Test collection**
 - Testing should be completed with Novel respiratory PPE (gown, gloves, N-95, eye protection) though is NOT considered an aerosol generating procedure (AGP)
 - Collect an Oropharyngeal/Nasopharyngeal or Mid-turbinate swab
- **Induction considerations**
 - Strongly recommend outpatient ripening for medical and elective inductions
- **Patient attire/location**
 - Recommend patient wear surgical mask throughout hospitalization
 - Limiting time outside of labor/antepartum rooms
- **Support/caregiver:**
 - Can switch support person every 24 hours during labor and postpartum admission
 - Required to wear facemask
 - Limit time in common areas including hallways
 - Handwashing often
 - If develops infectious symptoms (cough, fevers, etc) MUST leave immediately
- **Labor management considerations**
 - Nitrous oxide can be used if COVID testing is negative
 - Only nasal cannula use for maternal hypoxia (NOT used for intrauterine resuscitation)
- **Provider PPE considerations**
 - While test is pending, providers should be wearing DROPLET PPE (surgical mask and eye protection)
 - If test is still pending at the time of 2nd stage, AGP sign should be placed on patient's door and providers should don Novel respiratory PPE (gown, gloves, N-95, eye protection)
 - Once test results as negative, all isolation orders should be removed and providers can return to universal precautions
 - Limit number of providers in the room for deliveries to conserve PPE
- **Considerations if asymptomatic patient undergoing a C-section**
 - If COVID negative, standard OR precautions should be applied for all team members
 - If COVID pending (for the asymptomatic patient), team members should be wearing Novel respiratory PPE (gown, gloves, N-95, eye protection) if patient is at high risk of requiring intubation/general anesthesia (see below)
- **Cases at High Risk for requiring general anesthesia:**
 - Epidural in situ > 12 hours
 - pain during labor requiring > or equal to 2 additional anesthesia administered boluses
 - non-CSE/DPE epidural catheters
 - concerns for morbidly adherent placenta
 - BMI > 40
 - emergent surgery
- **Strong recommendation for early PP discharge if mom and baby are stable**
 - If patient is re-admitted and COVID negative, caregiver and newborn can also come to the hospital with the patient
- **House cleaning considerations**
 - Do not need to delay room cleaning/turnover after asymptomatic COVID test collection (not an AGP)
 - Need to wait 1 hour prior to room cleaning after AGP (ie. 2nd stage or C-section with intubation) for COVID unknown/pending patient

Inpatient Antepartum, Labor and Delivery, Postpartum Guidelines
COVID 19 (PUI or CONFIRMED)
HANDLING OF SPECIMENS FOR LAB AND PATHOLOGY

CONTACTING PATHOLOGY

- Normal Business hours M-F 7am-6pm
 - Contact the Mission Bay Gross Room: 514-3711
 - Provide patient's name and Medical Record Number (MRN)
 - Notify verbally of COVID status when to expect specimen to be brought to Gross Room (M2379)
- After-Hours (Weekends)
 - Please contact on-call resident pager: 443-1166
 - Provide the patient's name and MRN
 - Notify verbally of suspected or confirmed COVID status and that specimen will be placed in Pathology pass-through refrigerator.
- Labeling of Requisition and Specimen Container
 - Both Requisition and Specimen container should be labeled with a SHARPIE with big letters:
 - Suspected COVID-19 or
 - Confirmed COVID-19
- Transporting of Specimen
 - DO NOT PLACE IN TUG
 - Specimens must be walked down to pathology gross room
 - During Normal Business hours – hand deliver to pathology gross room (M2379 – 2nd floor of Gateway Medical Building, next to lab/bloodbank)
 - After-hours – hand deliver to pathology gross room pass-through refrigerator (M2379, badge access to the door that leads to refrigerator)

Labs (including cord gases)

- Ensure proper labeling of specimen (specimen label matches lab requisition, double RN check)
- Place contaminated lab biohazard bag with specimen in a new clean lab biohazard bag (double bag)
- Lab requisition to be placed in the outside pocket of lab bag
- These CAN be sent via pneumatic tube (ensure specimens are tightly sealed)
- Labeling of PUI or COVID + on specimen or requisition is not necessary